Muscle Biopsy Procedure

Richard J. Piercy MA VetMB MS PhD Dip.ACVIM MRCVS

Comparative Neuromuscular Diseases Laboratory Royal Veterinary College Royal College Street London NW1 0TU

Email: neuromuscular@rvc.ac.uk

Please contact the laboratory W $\mathbb{R} \mathbb{R} \mathbb{R} \mathbb{N} \setminus \mathbb{R} \otimes \mathbb{V} \mathbb{D} \mathbb{P} \mathbb{S} O \mathbb{H} \mathbb{B} \mathbb{P}$ Dals Over can only accept pre-booked samples. Samples must be pre-booked at least K R X U V before submitting, however we cannot guarantee to have spaces available within th short time frame. If this is the case we will offer alternative dates for submission.

Indications / possible diagnoses Exertional myopathies (RER/EPSM etc) Materials re quired

Muscle Biopsy Method

- 1. Contact the laboratory WR SUH ERRN \RXU VDPSOHV DW OHDV 2rganise same day or overnight courier service prior to sampling.
- 2. For horses with exertional myopathies the semimembranosis is generally chosen; for equine motor neuron disease, biopsy the sacrocaudalis dorsalis (craniolateral to tail head). For other disorders, choose the site based on the most obviously affected muscle or biopsy several regions.
- 3. Sedate horse and prepare skin for sterile surgery.
- 4. A Bergstrom biopsy needle is suitable, but reliable (and usually better) results are obtained with open biopsy.
- 5. Inject subcutaneously up to 10 ml of local anaesthetic, taking care to avoid direct injection into the muscle layer.
- 6.